

## Math 8062 Homework 3

Due Wednesday, 2/14/18

**1.** There is a standard way to glue together two (connected) manifolds  $M$  and  $N$  of the same dimension. Remove an open ball  $B^n$  from each of  $M$  and  $N$ , and glue  $M \setminus B^n$  to  $N \setminus B^n$  along the two  $(n - 1)$  dimensional boundary spheres. The resulting manifold is called the *connected sum* of  $M$  and  $N$ , and is denoted  $M \# N$ .

a) Prove that when  $n \geq 3$ ,  $\pi_1(M \setminus B^n) \cong \pi_1(M)$ .

b) Prove that when  $n \geq 3$ ,  $\pi_1(M \# N) \cong \pi_1(M) * \pi_1(N)$ .

**2.** Let  $S$  be a compact, connected surface with boundary. Let  $X = X^2$  be a cell complex structure on  $S$ , in which every 2-cell is an embedded polygon. Prove that  $X$  deformation retracts into the 1-skeleton  $X^1$ , and conclude that  $\pi_1(S)$  is a free group.

For extra credit, extend this argument to dimension  $n$ : that is, show an  $n$ -manifold with boundary deformation retracts into its  $(n - 1)$  skeleton.

**3.** Problem 8 on page 53 of Hatcher.

**4.** Problem 14 on page 54 of Hatcher. For extra credit, show that  $X$  is a 3-manifold.

*General hint:* for some of these problems, van Kampen's theorem will be more useful than Proposition 1.26 (the application to cell complexes).